

The nonminimal scalar multiplet coupled to supersymmetric Yang-Mills

Silvia Penati¹ and Daniela Zanon²

Dipartimento di Fisica dell'Università di Milano and
INFN, Sezione di Milano, via Celoria 16, I-20133 Milano, Italy

ABSTRACT

We consider the coupling of nonminimal scalar multiplets to supersymmetric Yang-Mills in four dimensions and compute the one-loop contribution to the low-energy effective action in the abelian sector. We show that the resulting theory realizes the dual version of the corresponding one from $N = 2$ supersymmetric Yang-Mills.

¹e-mail: penati@mi.infn.it

²e-mail: zanon@mi.infn.it

Recently much attention has been given to the study of low-energy effective actions for supersymmetric gauge theories in four dimensions. These investigations were boosted primarily by the Seiberg and Witten's exact construction [1] of the abelian low-energy effective action for the $N = 2$ Yang-Mills theory. In a $N = 1$ superspace formalism such a theory is described in terms of chiral superfields, in the adjoint representation of the gauge group, minimally coupled to the vector gauge multiplet.

In this paper we consider the corresponding situation where the chiral superfields are replaced by complex linear superfields [2], [3]. It is well known that the on-shell spectrum of the minimal scalar, described by a chiral superfield, and of the nonminimal one, described by a complex linear superfield, are the same, the two differing only in their auxiliary field content. The kinetic action for a complex linear superfield Σ , $\bar{\Sigma}$, satisfying $\bar{D}^2\Sigma = 0$, $D^2\bar{\Sigma} = 0$ is

$$S_{kin} = - \int d^4x d^4\theta \bar{\Sigma}\Sigma \quad (1)$$

In terms of component fields [2], [3], it becomes (the minus sign in (1) has been chosen so that it gives the correct sign for the kinetic term of the component scalar field)

$$S = \int d^4x [\bar{B}\square B - \bar{\zeta}^{\dot{\alpha}}i\partial_{\alpha\dot{\alpha}}\zeta^{\alpha} - \bar{H}H + \beta^{\alpha}\rho_{\alpha} + \bar{\beta}^{\dot{\alpha}}\bar{\rho}_{\dot{\alpha}} + \bar{p}^{\alpha\dot{\alpha}}p_{\alpha\dot{\alpha}}] \quad (2)$$

with physical components $B = \Sigma|$, $\zeta_{\alpha} = D_{\alpha}\bar{\Sigma}|$ like in the chiral superfield, but different auxiliary modes $\rho_{\alpha} = D_{\alpha}\Sigma|$, $H = D^2\Sigma|$, $p_{\alpha\dot{\alpha}} = \bar{D}_{\dot{\alpha}}D_{\alpha}\Sigma|$, $\bar{\beta}_{\dot{\alpha}} = \frac{1}{2}D^{\alpha}\bar{D}_{\dot{\alpha}}D_{\alpha}\Sigma|$.

The coupling of Σ to the Yang-Mills superfield V is easily accomplished by defining covariant derivatives and covariantly linear superfields with respect to the gauge fields. We use the conventions in ref. [3]. Thus we consider $N = 1$ superfields V , Σ and $\bar{\Sigma}$ Lie-algebra valued in the adjoint representation, $V = V^a T_a$, $\Sigma = \Sigma^a T_a$, $\bar{\Sigma} = \bar{\Sigma}^a T_a$ with $[T_a, T_b] = if_{ab}^c T_c$ and $tr T_a T_b = K\delta_{ab}$, and we introduce covariant derivatives, in vector representation,

$$\begin{aligned} \nabla_{\alpha} &= e^{-\frac{V}{2}} D_{\alpha} e^{\frac{V}{2}} & \bar{\nabla}_{\dot{\alpha}} &= e^{\frac{V}{2}} \bar{D}_{\dot{\alpha}} e^{-\frac{V}{2}} \\ \nabla_{\alpha\dot{\alpha}} &= -i\{\nabla_{\alpha}, \bar{\nabla}_{\dot{\alpha}}\} \end{aligned} \quad (3)$$

We define the Yang-Mills field-strength

$$W_{\alpha} = \frac{i}{2}[\bar{\nabla}^{\dot{\alpha}}, \{\bar{\nabla}_{\dot{\alpha}}, \nabla_{\alpha}\}] \quad (4)$$

and covariantly linear superfields $\Sigma_c, \bar{\Sigma}_c$ subject to the constraints $\bar{\nabla}^2 \Sigma_c = 0, \nabla^2 \bar{\Sigma}_c = 0$. In vector representation they are given by

$$\Sigma_c = e^{\frac{V}{2}} \Sigma e^{-\frac{V}{2}} \quad \bar{\Sigma}_c = e^{-\frac{V}{2}} \bar{\Sigma} e^{\frac{V}{2}} \quad (5)$$

The gauge invariant classical action is

$$S = \frac{1}{\beta^2} \left[\int d^4x d^2\theta \frac{1}{4} \text{tr}(W^\alpha W_\alpha) + \int d^4x d^2\bar{\theta} \frac{1}{4} \text{tr}(\bar{W}^{\dot{\alpha}} \bar{W}_{\dot{\alpha}}) - \int d^4x d^2\theta d^2\bar{\theta} \text{tr}(\bar{\Sigma}_c \Sigma_c) \right] \quad (6)$$

In terms of the prepotential V and of the complex linear superfields Σ 's, with gauge transformations

$$\begin{aligned} e^{V'} &= e^{i\bar{\Lambda}} e^V e^{-i\Lambda} & e^{-V'} &= e^{i\Lambda} e^{-V} e^{-i\bar{\Lambda}} & \bar{D}_{\dot{\alpha}} \Lambda &= 0, & D_{\alpha} \bar{\Lambda} &= 0 \\ \Sigma' &= e^{i\Lambda} \Sigma e^{-i\Lambda} & \bar{\Sigma}' &= e^{i\bar{\Lambda}} \bar{\Sigma} e^{-i\bar{\Lambda}} \end{aligned} \quad (7)$$

it can be rewritten as

$$S = \frac{1}{\beta^2} \int d^4x d^2\theta d^2\bar{\theta} \text{tr} \left[-\frac{1}{2} (e^{-V} D^\alpha e^V) \bar{D}^2 (e^{-V} D_\alpha e^V) - e^{-V} \bar{\Sigma} e^V \Sigma \right] \quad (8)$$

In addition to the explicit $N = 1$ supersymmetry, the action in (6) possesses an invariance under a second set of global supersymmetry transformations

$$\begin{aligned} \delta \bar{\Sigma}_c &= -i\epsilon^\alpha W_\alpha & \delta \Sigma_c &= i\bar{\epsilon}^{\dot{\alpha}} \bar{W}_{\dot{\alpha}} \\ \delta \nabla_\alpha &= \epsilon_\alpha \Sigma_c & \delta \bar{\nabla}_{\dot{\alpha}} &= -\bar{\epsilon}_{\dot{\alpha}} \bar{\Sigma}_c \end{aligned} \quad (9)$$

The corresponding variations of the field-strengths can be obtained using the definitions in (4)

$$\delta W_\alpha = -\bar{\epsilon}^{\dot{\alpha}} \nabla_{\alpha\dot{\alpha}} \bar{\Sigma}_c - \frac{i}{2} \bar{\epsilon}^{\dot{\alpha}} \nabla_\alpha \bar{\nabla}_{\dot{\alpha}} \bar{\Sigma}_c \quad \delta \bar{W}_{\dot{\alpha}} = \epsilon^\alpha \nabla_{\alpha\dot{\alpha}} \Sigma_c + \frac{i}{2} \epsilon^\alpha \bar{\nabla}_{\dot{\alpha}} \nabla_\alpha \Sigma_c \quad (10)$$

The supersymmetry algebra in (9) and (10) closes on-shell. Indeed the commutators of two such transformations give

$$\begin{aligned} [\delta_1, \delta_2] \bar{\Sigma}_c &= (\epsilon_2^\alpha \bar{\epsilon}_1^{\dot{\alpha}} - \epsilon_1^\alpha \bar{\epsilon}_2^{\dot{\alpha}}) (i\nabla_{\alpha\dot{\alpha}} - \frac{1}{2} \nabla_\alpha \bar{\nabla}_{\dot{\alpha}}) \bar{\Sigma}_c \\ [\delta_1, \delta_2] \Sigma_c &= (\bar{\epsilon}_2^{\dot{\alpha}} \epsilon_1^\alpha - \bar{\epsilon}_1^{\dot{\alpha}} \epsilon_2^\alpha) (i\nabla_{\alpha\dot{\alpha}} - \frac{1}{2} \bar{\nabla}_{\dot{\alpha}} \nabla_\alpha) \Sigma_c \\ [\delta_1, \delta_2] W_\alpha &= (\bar{\epsilon}_2^{\dot{\alpha}} \epsilon_1^\beta - \bar{\epsilon}_1^{\dot{\alpha}} \epsilon_2^\beta) (i\nabla_{\beta\dot{\alpha}} W_\alpha + c_{\beta\alpha} \bar{\nabla}_{\dot{\alpha}} \nabla^\gamma W_\gamma) \\ [\delta_1, \delta_2] \bar{W}_{\dot{\alpha}} &= (\epsilon_2^\alpha \bar{\epsilon}_1^{\dot{\beta}} - \epsilon_1^\alpha \bar{\epsilon}_2^{\dot{\beta}}) (i\nabla_{\alpha\dot{\beta}} \bar{W}_{\dot{\alpha}} + c_{\beta\dot{\alpha}} \nabla_\alpha \bar{\nabla}^{\dot{\gamma}} \bar{W}_{\dot{\gamma}}) \end{aligned} \quad (11)$$

and the closure of the algebra is implemented once the equations of motion, $\bar{\nabla}_{\dot{\alpha}}\bar{\Sigma}_c = 0$, $\nabla_{\alpha}\Sigma_c = 0$, $\nabla^{\alpha}W_{\alpha} = 0$, $\bar{\nabla}^{\dot{\alpha}}\bar{W}_{\dot{\alpha}} = 0$, are imposed. The $N = 1$ description in terms of complex linear superfields and gauge multiplets is missing auxiliary degrees of freedom needed for the off-shell closure of the $N = 2$ supersymmetry algebra. This is in contradistinction with the corresponding situation for covariantly chiral superfields coupled to Yang-Mills: there the second supersymmetry which realizes the $N = 2$ invariance closes off-shell. This statement can be easily checked considering the $N = 2$ Yang-Mills action written in terms of $N = 1$ superfields

$$S = \frac{1}{g^2} \left[\int d^4x d^2\theta \frac{1}{4} \text{tr}(\mathcal{W}^{\alpha}\mathcal{W}_{\alpha}) + \int d^4x d^2\bar{\theta} \frac{1}{4} \text{tr}(\bar{\mathcal{W}}^{\dot{\alpha}}\bar{\mathcal{W}}_{\dot{\alpha}}) + \int d^4x d^2\theta d^2\bar{\theta} \text{tr}(\bar{\Phi}_c\Phi_c) \right] \quad (12)$$

where Φ_c , $\bar{\Phi}_c$ are covariantly chiral, antichiral superfields respectively, and $\mathcal{W}_{\alpha}$ is the Yang-Mills field-strength. The action in (12) is invariant under

$$\begin{aligned} \delta\Phi_c &= -i\xi^{\alpha}\mathcal{W}_{\alpha} & \delta\bar{\Phi}_c &= i\bar{\xi}^{\dot{\alpha}}\bar{\mathcal{W}}_{\dot{\alpha}} \\ \delta\nabla_{\alpha} &= \xi_{\alpha}\bar{\Phi}_c & \delta\bar{\nabla}_{\dot{\alpha}} &= -\bar{\xi}_{\dot{\alpha}}\Phi_c \end{aligned} \quad (13)$$

with corresponding transformations for the field-strengths

$$\delta\mathcal{W}_{\alpha} = i\xi_{\alpha}\bar{\nabla}^2\bar{\Phi}_c - \bar{\xi}^{\dot{\alpha}}\nabla_{\alpha\dot{\alpha}}\Phi_c \quad \delta\bar{\mathcal{W}}_{\dot{\alpha}} = -i\bar{\xi}_{\dot{\alpha}}\nabla^2\Phi_c + \xi^{\alpha}\nabla_{\alpha\dot{\alpha}}\bar{\Phi}_c \quad (14)$$

Using (13) and (14) it is straightforward to compute the commutator algebra and verify the off-shell closure.

We now go back to the action in (8) and consider its quantization. We shall restrict our attention to the $SU(2)$ case, with $f_{abc} = \epsilon_{abc}$ and $(T_a)_{kl} = i\epsilon_{kal}$. The aim is to perform the following one-loop computation: we give a vacuum expectation value to the complex linear superfields in an abelian direction

$$\Sigma = (0, 0, \Sigma) \quad \bar{\Sigma} = (0, 0, \bar{\Sigma}) \quad (15)$$

Then we compute all one-loop diagrams with external Σ , $\bar{\Sigma}$ and no spinor nor space-time derivatives acting on them. Obviously this amounts to the determination of the leading contribution to the low-energy effective action, exactly as in the corresponding calculation for the $N = 2$ supersymmetric Yang-Mills theory.

In order to proceed in the perturbative calculation, it is convenient to make a quantum-background splitting, $\Sigma \rightarrow \Sigma_Q + \bar{\Sigma}$, $\bar{\Sigma} \rightarrow \bar{\Sigma}_Q + \bar{\Sigma}$, so that Σ and $\bar{\Sigma}$ are the external fields given in (15), while Σ_Q , $\bar{\Sigma}_Q$ and V are the quantum fields. Since one does not know how to perform superspace functional differentiation and integration on complex linear superfields, a direct quantization is not available. Here we follow the approach proposed in some recent papers [4], [5]: the linearity constraints are solved in terms of unconstrained gauge spinor superfields σ^α , $\bar{\sigma}^{\dot{\alpha}}$ whose field-strengths are $\Sigma_Q = \bar{D}_{\dot{\alpha}} \bar{\sigma}^{\dot{\alpha}}$, $\bar{\Sigma}_Q = D_\alpha \sigma^\alpha$. The gauge invariance introduced in this manner needs gauge fixing; in fact an infinite set of invariances arises and it leads to a tower of ghosts. In ref. [4] it was shown how to proceed systematically in the quantization using the Batalin-Vilkovisky method. In ref. [5] this formulation has been applied to the calculation of the one-loop β -function for the nonlinear σ -model defined in terms of nonminimal scalar multiplets. It has been shown that choosing gauge-fixing functions independent of the external background all the ghosts essentially decouple; the calculation becomes manageable and no formal (infinite) manipulations are necessary.

As compared to the nonlinear σ -model case, we have to face here the extra requirement of maintaining the gauge invariance with respect to the Yang-Mills fields. This is most easily done in a covariant approach: following the gauge-fixing procedure described in detail in ref. [4], the only thing we need to do is to replace the flat derivatives with the corresponding covariant ones given in (3). This automatically leads to the coupling of the infinite tower of ghosts to the Yang-Mills fields. While in general this might be quite difficult to handle, no extra complications arise for the calculation we have in mind. In fact it is immediate to realize that these ghosts never couple to the external background Σ , $\bar{\Sigma}$ and then they do not contribute at one loop.

In ref. [5] the $\langle \sigma^\alpha \bar{\sigma}^{\dot{\alpha}} \rangle$ propagator has been computed and we refer the reader to that paper for the details of the calculation. The final result is

$$\begin{aligned} \langle \sigma_\alpha^a \bar{\sigma}_{\dot{\alpha}}^b \rangle = & \frac{\beta^2}{K} \delta^{ab} \left[-\frac{i\partial_{\alpha\dot{\alpha}}}{\square} + \frac{3(kk'_1)^2 + 4 - 2k_1'^2}{4(kk'_1)^2} i\partial_{\alpha\dot{\alpha}} \frac{D^2 \bar{D}^2}{\square^2} + \right. \\ & \left. + \frac{3k^2 - 2}{4k^2} i\partial_{\alpha\dot{\alpha}} \frac{D_\beta \bar{D}^2 D^\beta}{\square^2} + \frac{2 - k^2}{4k^2} i\partial_{\alpha\dot{\beta}} i\partial_{\beta\dot{\alpha}} \frac{D^\beta \bar{D}^{\dot{\beta}}}{\square^2} \right] \delta^{(4)}(\theta - \theta') \quad (16) \end{aligned}$$

where k and k'_1 are gauge parameters. The gauge fixing for the $N = 1$ vector multiplet is performed in standard manner: as in refs. [6] and [7] we choose the supersymmetric Landau gauge which gives the V -propagator

$$\langle V^a V^b \rangle = -\frac{\beta^2}{K} \delta^{ab} \frac{D^\alpha \bar{D}^2 D_\alpha}{\square^2} \delta^{(4)}(\theta - \theta') \quad (17)$$

Again the chiral ghosts introduced by the quantization in the Yang-Mills sector do not couple to the external fields and therefore do not enter a one-loop calculation.

After quantum-background splitting, from the action in (8) we have as relevant interactions the cubic vertices

$$V^{(3)} = -\frac{1}{\beta^2} \text{tr}(\bar{\Sigma}[V, \Sigma_Q]) - \frac{1}{\beta^2} \text{tr}([\bar{\Sigma}_Q, V]\Sigma) \quad (18)$$

and the quartic vertex

$$V^{(4)} = -\frac{1}{2\beta^2} \text{tr}(\bar{\Sigma}[V, [V, \Sigma]]) \quad (19)$$

First we notice that, since the quantum fields σ_α , $\bar{\sigma}_{\dot{\alpha}}$ always interact with the external background through their field-strengths Σ_Q , $\bar{\Sigma}_Q$, instead of the propagator in (16) it is sufficient to use the much simpler expression [5]

$$\langle \bar{\Sigma}_Q^a \Sigma_Q^b \rangle = D^\alpha \langle \sigma_\alpha^a \bar{\sigma}_{\dot{\alpha}}^b \rangle \bar{D}^{\dot{\alpha}} = \frac{\beta^2}{K} \delta^{ab} \left[\frac{D^2 \bar{D}^2}{\square} + \frac{D_\alpha \bar{D}^2 D^\alpha}{\square} \right] \delta^{(4)}(\theta - \theta') \quad (20)$$

Then we find convenient to consider effective V -propagators which contain the sum of all possible insertions of the quartic interaction in (19). The sum is performed taking advantage of the fact that, since we want to compute the leading contributions to the effective action, no derivative acts on the external fields. For the spinor derivatives present in (17) we use the relation $(D^\alpha \bar{D}^2 D_\alpha)^n = (-\square)^{n-1} D^\alpha \bar{D}^2 D_\alpha$ and obtain

$$\langle\langle V^a V^b \rangle\rangle = -\frac{\beta^2}{K} \begin{pmatrix} (\square + \bar{\Sigma}\Sigma)^{-1} & 0 & 0 \\ 0 & (\square + \bar{\Sigma}\Sigma)^{-1} & 0 \\ 0 & 0 & \square^{-1} \end{pmatrix}^{ab} \frac{D^\alpha \bar{D}^2 D_\alpha}{\square} \delta^{(4)}(\theta - \theta') \quad (21)$$

Now the one-loop $\Sigma\bar{\Sigma}$ low-energy effective action is easily computed, summing over graphs with n three-point vertices $\frac{-1}{\beta^2} \text{tr}(\bar{\Sigma}[V, \Sigma_Q])$ and n vertices $\frac{-1}{\beta^2} \text{tr}([\bar{\Sigma}_Q, V]\Sigma)$, with n propagators $\langle\langle V^a V^b \rangle\rangle$ and n propagators $\langle \bar{\Sigma}_Q^a \Sigma_Q^b \rangle$. The D -algebra is straightforward. We have spinor derivatives from the propagators in (20) and (21). Since they all have to stay in the loop, we repeatedly use the identities

$$D^\alpha \bar{D}^2 D_\alpha (D^2 \bar{D}^2 + D_\beta \bar{D}^2 D^\beta) = D^\alpha \bar{D}^2 D_\alpha (\bar{D}^2 D^2 + \bar{D}_{\dot{\beta}} D^2 \bar{D}^{\dot{\beta}}) = \square D^\alpha \bar{D}^2 D_\alpha \quad (22)$$

to reduce the number of D 's and $\bar{D}$'s. As a final step the D -algebra is completed by making use of

$$\delta^{(4)}(\theta - \theta') D^2 \bar{D}^2 \delta^{(4)}(\theta - \theta') = \frac{1}{2} \delta^{(4)}(\theta - \theta') D^\alpha \bar{D}^2 D_\alpha \delta^{(4)}(\theta - \theta') = \delta^{(4)}(\theta - \theta') \quad (23)$$

The total sum of these contributions gives

$$\begin{aligned}\Gamma_{(1)} &= \int \frac{d^4p}{(2\pi)^4} d^4\theta \frac{2}{p^2} \sum_{n=1}^{\infty} \frac{1}{n} \left(2 \frac{-\bar{\Sigma}\Sigma}{p^2 - \bar{\Sigma}\Sigma} \right)^n \\ &= - \int \frac{d^4p}{(2\pi)^4} d^4\theta \frac{2}{p^2} \left[\ln\left(1 + \frac{\Sigma\bar{\Sigma}}{p^2}\right) - \ln\left(1 - \frac{\Sigma\bar{\Sigma}}{p^2}\right) \right]\end{aligned}\quad (24)$$

In addition one has the contribution from the effective V -propagator in (21) with no extra background insertions: it gives

$$\Gamma_{(2)} = - \int \frac{d^4p}{(2\pi)^4} d^4\theta \frac{2}{p^2} \ln\left(1 - \frac{\Sigma\bar{\Sigma}}{p^2}\right) \quad (25)$$

Summing the terms in (24) and (25) one obtains

$$\Gamma = - \frac{2}{(4\pi)^2} \int_0^\infty dp^2 d^4\theta \ln\left(1 + \frac{\Sigma\bar{\Sigma}}{p^2}\right) \quad (26)$$

The integral is divergent: it gives rise to a wavefunction renormalization of the Σ 's plus a finite contribution

$$\Gamma_{eff}(\bar{\Sigma}, \Sigma) = \frac{-2}{(4\pi)^2} \bar{\Sigma}\Sigma \ln\left(\frac{\bar{\Sigma}\Sigma}{\mu^2}\right) \quad (27)$$

where μ is a renormalization mass. The result in (27) represents the one-loop low-energy contribution to the $\bar{\Sigma}\Sigma$ lagrangian, exactly as in the corresponding calculation in terms of chiral superfields. Now we elaborate on this.

The equivalence of the descriptions of the scalar multiplet by the complex linear superfield and by the chiral superfield can be understood in terms of a duality transformation. One writes a first order action

$$S = \int d^4x d^4\theta \left[\bar{\Phi}\Phi + \Sigma\Phi + \bar{\Sigma}\bar{\Phi} \right] \quad (28)$$

with Σ satisfying the constraint $\bar{D}^2\Sigma = 0$, (which implies $\Sigma = \bar{D}_{\dot{\alpha}}\bar{\sigma}^{\dot{\alpha}}$), and Φ unconstrained. Using the equations of motion to eliminate Φ , $\bar{\Phi}$, one obtains the linear superfield action. Eliminating instead Σ , $\bar{\Sigma}$ whose equations of motion impose the chirality constraint on Φ ($\bar{\Phi}$), $\bar{D}_{\dot{\alpha}}\Phi = 0$, ($D_{\alpha}\bar{\Phi} = 0$), one obtains the standard chiral superfield action. In view of this observation our one-loop calculation has shown that this classical duality of the matter fields is maintained at the quantum level. Since we have shown that

both theories in (6) and in (12) are invariant under a second supersymmetry which mixes the matter fields with the gauge fields, we would expect the duality in the matter sector, $\Phi_c \rightarrow \Sigma_c$, to be accompanied by the electro-magnetic duality, $\mathcal{W}_\alpha \rightarrow W_\alpha$, in the vector sector.

However things do not quite work so simply. We can start for example with the linear multiplet Σ_c , satisfying the constraint $\bar{\nabla}^2 \Sigma_c = 0$ (which implies $\Sigma_c = \bar{\nabla}_{\dot{\alpha}} \bar{\sigma}^{\dot{\alpha}}$) and the gauge field strength $W_\alpha = \frac{i}{2}[\bar{\nabla}^{\dot{\alpha}}, \{\bar{\nabla}_{\dot{\alpha}}, \nabla_\alpha\}]$, and consider the first order action

$$\begin{aligned} S = & \frac{1}{g^2} tr \left[\int d^4x d^4\theta \bar{\Phi}_c \Phi_c + \frac{1}{4} \int d^4x d^2\theta \mathcal{W}^\alpha \mathcal{W}_\alpha + \frac{1}{4} \int d^4x d^2\bar{\theta} \bar{\mathcal{W}}^{\dot{\alpha}} \bar{\mathcal{W}}_{\dot{\alpha}} \right] \\ & + tr \int d^4x d^4\theta (\Phi_c \Sigma_c + \bar{\Phi}_c \bar{\Sigma}_c) + \frac{i}{2} tr \int d^4x d^2\theta \mathcal{W}^\alpha W_\alpha \\ & - \frac{i}{2} tr \int d^4x d^2\bar{\theta} \bar{\mathcal{W}}^{\dot{\alpha}} \bar{W}_{\dot{\alpha}} \end{aligned} \quad (29)$$

with unconstrained superfields $\Phi_c, \bar{\Phi}_c$, and chiral spinors $\mathcal{W}_\alpha$ and $\bar{\mathcal{W}}_{\dot{\alpha}}$.

Now, functional integration over $\Phi_c, \bar{\Phi}_c, \mathcal{W}_\alpha$ and $\bar{\mathcal{W}}_{\dot{\alpha}}$, leads to the action

$$S = g^2 tr \left[- \int d^4x d^4\theta \bar{\Sigma}_c \Sigma_c + \frac{1}{4} \int d^4x d^2\theta W^\alpha W_\alpha + \frac{1}{4} \int d^4x d^2\bar{\theta} \bar{W}^{\dot{\alpha}} \bar{W}_{\dot{\alpha}} \right] \quad (30)$$

which exactly matches the one in eq. (6) with the identification $\beta^2 = 1/g^2$. On the other hand, using the equations of motion to eliminate the superfields $\Sigma, \bar{\Sigma}$ and V , we obtain the chirality constraints

$$\bar{\nabla}_{\dot{\alpha}} \Phi_c = 0 \quad \nabla_\alpha \bar{\Phi}_c = 0 \quad (31)$$

but *not* the Bianchi identities

$$\nabla^\alpha \mathcal{W}_\alpha + \bar{\nabla}^{\dot{\alpha}} \bar{\mathcal{W}}_{\dot{\alpha}} = 0 \quad (32)$$

Instead of eq. (32) one obtains the equations of motion

$$\nabla^\alpha \mathcal{W}_\alpha = 0 \quad \bar{\nabla}^{\dot{\alpha}} \bar{\mathcal{W}}_{\dot{\alpha}} = 0 \quad (33)$$

so that in a certain sense the standard $N = 2$ Yang-Mills theory is recovered only on-shell. We believe that this is just a consequence of the fact that for the theory in (6) the second supersymmetry is an on-shell invariance.

We notice that the first order action in (29) is appropriate for describing the above mentioned duality equivalence in the abelian case. Indeed for the abelian theory we delete the tr operation and consider the linear multiplet satisfying the constraint $\bar{D}^2 \Sigma = 0$ and the gauge field strength $W_\alpha = i\bar{D}^2 D_\alpha V$. Using the field equations to eliminate $\Sigma, \bar{\Sigma}$ and

V , one recovers the chirality conditions on Φ and $\bar{\Phi}$ and the Bianchi identities on $\mathcal{W}_\alpha$, $\bar{\mathcal{W}}_{\dot{\alpha}}$, so that the abelian $N = 2$ Yang-Mills theory is reconstructed.

In the nonabelian case one should try to obtain a formulation of the theory in terms of complex linear superfields and gauge fields directly in a $N = 2$ description. Indeed in ref. [1] the complete, nonperturbative information about the low-energy effective action was obtained starting from the $N = 2$ superspace formulation of the theory in (12). In $N = 2$ superspace the action is given by the chiral integral of the prepotential, a holomorphic function $\mathcal{F}(W)$ being W the $N = 2$ gauge superfield strength [8]. For the action in (30) such a formulation is not known. In order to move into $N = 2$ superspace, first one should learn which are the auxiliary degrees of freedom not present in the $N = 1$ formulation of the theory. Second one should find a $N = 2$ superfield which might accomodate a complex linear superfield and a Yang-Mills field-strength among its $N = 1$ components. These issues seem worth to pursue.

In a different but complementary perspective, it could be interesting to consider models with mixed matter, i.e. models constructed in terms of chiral and complex linear superfields along the lines studied in [9], and analyze their interaction with the Yang-Mills gauge multiplet.

As a final comment we observe that in order to have these systems completely under control it would be necessary to understand how to treat the infinite tower of ghosts introduced by the Batalin-Vilkovisky quantization of the complex linear superfield. From the results presented here it is clear that the contribution to the one-loop low-energy effective action of the gauge field could be obtained using $N = 2$ supersymmetry. This indicates that the final answer is simple, and therefore we only need an intelligent way to handle efficiently the ghost calculation.

Acknowledgments

We would like to thank M. Grisaru for useful discussions. This work was supported in part by MURST and by the European Commission TMR programme ERBFMRX-CT96-0045, in which the authors are associated to the University of Torino.

References

- [1] N. Seiberg and E. Witten, Nucl. Phys. **B426** (1994) 19.
- [2] S.J. Gates and W. Siegel, Nucl. Phys. **B187** (1981) 389.
- [3] S.J. Gates, M.T. Grisaru, M. Roček and W. Siegel, *Superspace* (Addison-Wesley, 1983)
- [4] M.T. Grisaru, A. Van Proeyen, D. Zanon, Nucl. Phys. **B502** (1997) 345.
- [5] S. Penati, A. Refolli, A. Van Proeyen, D. Zanon, “*The nonminimal scalar multiplet: duality, sigma-model, beta-function*”, IFUM-579-FT, KUL-TF-97/26, hep-th/9710166.
- [6] B. de Wit, M.T. Grisaru, M. Roček, Phys. Lett. **B374** (1996) 297.
- [7] A. De Giovanni, M.T. Grisaru, M. Roček, R. von Unge, D. Zanon, Phys. Lett. **B409** (1997) 251.
- [8] G. Sierra and P.K. Townsend, in ”Supersymmetry and Supergravity 1983”, proc. XIXth Winter School, Karpacz, ed. B. Milewski (World Scientific, 1983), p. 396;
B. de Wit, P.G. Lauwers, R. Philippe, S.-Q. Su and A. Van Proeyen, Phys. Lett. **134B** (1984) 37;
S.J. Gates, Nucl. Phys. **B238** (1984) 349
- [9] S.J. Gates, Phys. Lett. **B365** (1996) 132, Nucl. Phys. **B485** (1997) 145;
S.J. Gates, M.T. Grisaru, M.E. Knutt-Wehlau, M. Roček and O.A. Soloviev, Phys. Lett. **B396** (1997) 167.